Immunohistochemical classification of breast tumours

Workshop in Diagnostic Immunohistochemistry Aalborg University Hospital NordiQC October 2nd - 4th 2019

Anne-Vibeke Lænkholm Department of Surgical Pathology, Zealand University Hospital, Slagelse Denmark



Agenda

- Immunohistochemical biomarkers for
 - Diagnostics
 - Benign Hyperplasia and Ductal Carcinoma in Situ
 - Ductal Carcinoma in Situ and Lobular Carcinoma in Situ
 - Carcinoma In Situ and Invasive Carcinoma
 - Histological subtype classification
 - Malignant breast tumors
 - Predictive/Prognostic markers
 - Estrogen Receptor
 - Progesteron Receptor
 - HER2
 - Ki67
 - PD-L1
 - Intrinsic subtype classification by surrogate IHC markers?
 - Tumor heterogeneity

Triple Test Diagnostic approach – Breast Tumours



ANATOMY OF BREAST

Modified apocrine sweat glands.

- Breast parenchyma → 12 to 20 lobes.
- Within each lobe Lactiferous duct

 branches repeatedly → leads to
 no. of terminal ducts → each leads
 to a lobule → contains multiple
 acini/alveoli → TDLU
 (TERMINAL DUCT + LOBULE)
- Spaces around the lobules and ducts and between the lobes are filled with fatty tissue, ligaments and connective tissue → STROMA



connective **Terminal duct lobular unit = TDLU** tissue duct lobule duct

Mammary gland epithelium Two types of epithelial cells are present: Luminal cells and myoepithelial cells





Myoepithelial cells with contractile function forming a meshwork that does not cover the entire basement membrane nor the entire luminal cell

Epithelial cells with specific immunohistochemical phenotype



*Smooth muscle myosin heavy chain

Benign hyperplasia Positive staining for myoepitelial cells



Ductal Carcinoma In Situ

CK14 Ductal Carcinoma In Situ



Monotonous epithelial proliferation within ducts

Invasive Carcinoma i.e. SMMHC*

present

Not present



Detecting "presence"

Detecting "absence"

* Smooth muscle myosin heavy chain, as detected with clone SMMS-1

Loss of E-Cadherin Lobular Carcinoma in situ Terminal duct lobular unit

E-cadherin: Cell Adhesion Molecule

Carcinoma in situ

- Ductal carcinoma in situ
 - 12-15% of malignant lesions in the Danish screening population Microcalcifications ٠

 - Risk of progression to invasive carcinoma

 - Surgery with free margins Radiation therapy after breast conserving surgery



- Lobular carcinoma in situ
 - Incidence 0.5 3.6%

 - Often incidental finding Multifocal and often bilateral
 - Slowly proliferating lesions Observation / screening



Breast cancer: Incidence and mortality Denmark

Bryst ASR (W), Kvinder alder 0-74



Invasive Breast Cancer Histological Subtypes

- Ductal : up to 80%
- Lobular: 5 14%
- Tubular: 2 8%
- Mucinous: 2 4 %
- Apocrine: 1 4%
- Papillary 1 2%
- Other





E-Cadherin Cell adhesion molecule

Loss of E-Cadherin in 90% of Invasive lobular Carcinoma



E-Cadherin positive Invasive Ductal Carcinoma



CDH1 (16q22.1) loss of function mutation or deletion resulting in loss of the adhesion molecule E-cadherin¹⁵

P120 catenin dislocated to the cytoplam in lobular carcinoma A supplement for classification of lobular neoplasia



Lobular carcinoma not candidate for neoadjuvant chemotherapy

Apocrine carcinoma classification

ΗE

Androgen Receptor



AR staining in IHC-basallike breast cancer as potential marker for AR targeted treatment



Prognostic and predictive biomarkers

HER2 positive breast cancer: 15% Family of four receptors in the HER family HER2: Growth factor tyrosine kinase receptor Mediate cell growth differentiation and survival



EGFR, epidermal growth factor receptor; HER, human epidermal growth factor

HER2 and Breast Cancer Progression



Science, Vol 235, 1987

Timeline of HER2 targeting

FDA approvals i Breast Cancer



The timeline demonstrates rapid and accelerated development of drugs (and indications). Abbreviation: FDA, U.S. Food and Drug Administration.

HER2 targeting: Traztuzumab, protocols initiated 2000-2001



HERA 11 years update

The relative risk of a disease-free survival event was reduced by 24%.

	A DFS bene (% follow-up time after selective crossover)	fit HR (95% CI)	DFS events: 1-year trastuzumab vs observation	B Median follow-up (% follow-up time after selective crossover)	Overall survival benefit	HR (95% CI)	Deaths: 1-year trastuzumab vs observation
	2005 1 year (0%)	0-54 (0-43-0-67)	127 vs 220	2005 1 year		0-76 (0-47-1-23)	29 vs 37
	2006 2 years (4-3%)	0-64 (0-54-0-76)	218 vs 321	2006 2 years (4-1%)		0-66 (0-47-0-91)	59 vs 90
	2008 4 years →	0-76 (0-66-0-87)	369 vs 458	2008 4 years (30-9%)		0-85 (0-70-1-04)	182 vs 213
	2012 8 years (48-6%)	0-76 (0-67-0-86)	471 vs 570	2012 8 years (45·5%)	•	0-76 (0-65-0-88)	278 vs 350
	2015 11 years (53-6%)	0-76 (0-68-0-86)	505 vs 608	2015 11 years (50-4%)	⊢ ⊣	0-74 (0-64-0-86	320 vs 405
ER			,	-	-		
nocitivo	<u>د</u>			D			
positive.	2005 1 year (0%)	0.60 (0.42-0.85)	53 vs 82	2005 1 year (0%)		1-67 (0-74-3-78)	16 vs 9
Absolute	2006 2 years (4-2%)	0-68 (0-51-0-89)	87 vs 123	2006 2 years (4-0%)		0-69 (0-39-1-23)	20 vs 29
benefit	2008 4 years (34-9%)	0-84 (0-68-1-03)	162 vs 188	2008 4 years (32-5%)		1-03 (0-75-1-42)	75 vs 75
10 yr DFS:	2012 8 years	0-81 (0-68-0-98)	218 vs 253	2012 8 years (47·2%) →		0-84 (0-66-1-06)	126 vs 146
5.6%	2015 11 years → ■ → →	0-80 (0-68-0-96)	236 vs 277	2015 11 years (51-9%)	-	0-81 (0-65-1-00)	148 vs 176
	E			F			
FR	2005 1 year (0%) • • • • •	0-50 (0-38-0-67)	74 vs 138	2005 1 year (0%)		0-47 (0-24-0-90)	13 vs 28
	2006 2 years	0-62	131 vs 198	2006 2 years		0-64	39 vs 61
Negative	2008 4 years	0.70	207 vs 270	2008 d years		0.75	107 vs 138
Absoluto	(32.5%)	(0.59-0.84)		(29-2%)	s - 6	(0.58-0.97)	
Absolute	2012 8 years (47-1%)	0-72 (0-61-0-84)	253 vs 317	2012 8 years (43-6%)		0-70 (0-56-0-86)	152 vs 204
benefit	2015 11 years (52.2%)	0-73 (0-62-0-85)	269 vs 331	2015 11 years (48.8%)		0-70 (0-57-0-85)	172 vs 229
10 vr DFS	5 i		3	6	1		2
	Favours trastuzumab	Favours obse	rvation	Favours trastuzuma	5 F	Favours observ	ation
8.0%							

A 6.5% absolute gain was found in overall survival at 11 years between those in the 1-year trastuzumab group versus those in the observation group.

Lancet. 2017 March 25; 389(10075): 1195-1205.

JOURNAL OF CLINICAL ONCOLOGY

Human Epidermal Growth Factor Receptor 2 Testing in Breast Cancer: American Society of Clinical Oncology/ College of American Pathologists Clinical Practice Guideline Focused Update



Two different assays

IHC is an assay at the single-cell level
 It will detect even an individual positive cell

- ISH is a population-based assay (mean number of HER2 gene copies evaluated by scoring 20 cells)
 - Dual probe (Ratio: HER2 gene copy numbers/CEN17)
 - Mono probe (HER2 gene copy numbers/cell)

HER2 IHC

HER2 3+ and ISH + : 14-15 % (DK)

HER2 FISH

Green: :centromere chromosome 17 Red : HER2 gene Dual probe: Amplified HER2/CEN17 ratio >

HER2 Gene/Protein Assay

HER2 amplified and HER2 IHC 3+

HER2 testing by validated dual-probe ISH assay

Lin, N. U. et al. J Clin Oncol; 26:798-805 2008

Relevance of measured ER and PR status on the effects of 5 years of tamoxifen on the 10 year probability of recurrence (EBCTCG) Lancet. 2011 August 27; 378(9793): 771–784.

Interpretation of ER IHC

ER positive 86% of breast carcinomas (DK) Cut off \geq 1% (regardless of intensity)

Allred method

Proportion Score (PS)	Observation	Intensity Score (IS)	Observation
0	NONE	0	None
1	1%	1	Weak
2	1-10%	2	Intermediate
3	10-33%	3	Strong
4	33-66%		-
5	66-100%		
Total Score			Interpretation
	Sum of proportion scor	e and intensity sc	ore
0-2			Negative
3-8			Positive

Interpretation of PgR IHC

Neoadjuvant treatment

- Neoadjuvant systemic therapy for early breast cancer.
 - Tumor down sizing / staging
 - pCR (pathological complete response) is an evaluable end point for determining the efficacy of the treatment.
 - Prognostic information (DFS)

HER2 IHC

Post treatment - surgery

Tumor characteristics and association with pCR Lobular carcinoma not recommended for neoadjuvant treatment

A		Percentage of patients achieving pathological complete response (95% Cl)	-
Clinical tumour stage			
T1 (n-785)		18-3 (15-7-21-2)	
T2 (n=7328)	+	19-9 (19-0-20-9)	
T3 (n=2493)	-+-	13-0 (11-7-14-3)	
T4a-c (n=781)		14-5 (12-1-17-1)	
T4d (n=482)	— <u>; </u>	16-0 (12-8-19-6)	
Clinical nodal status			
Negative (n=6320)	+	18-8 (17-9-19-8)	
Positive (n= 5487)	+	16-9 (15-9-17-9)	
Histological type			
Ductal (n=8567)		15-5 (147-16-3)	
Lobular (n=1221)		7-8 (0-3-9-4)	pCR: 7.8%
Turnour anada		113 (130-100)	
1 (n= 426)		7864107	
2 (n=4307)	· +	12.3 (11.3, 13.3)	
2 (n=3217)	·	25.8 (24.3, 27.4)	
Clinical tumour subtype		*7.0 (mt.2.1.4)	
Hormone-receptor-positive. HER2-negative. grade 1/2 (n=1986)	+	7.5 (6-3-8-7)	
Hormone-receptor-positive, HER2-negative, grade 3 (n= 630)		16-2 (13-4-19-3)	
HER2- positive, hormone-receptor-positive, trastuzumab (n= 385)		30-9 (26-3-35-8)	
HER2-positive, hormone-receptor-positive, no trastuzumab (n=701)	— — — —	18-3 (15-5-21-3)	
HER2-positive, hormone-receptor-negative, trastuzumab (n=364)		50-3 (45-0-55-5) n	CR · 50.3%
HER2 positive, hormone-receptor-negative, no traster unab (n=471)		<u> 30-2 (26-0-34-5)</u>	
Triple negative (n= 1157)		33-6 (30-9-36-4)	
	0 10 20 30 40 Pathological complete response	50 60 (%)	
В		HR (95% CI)	

Cortazar et al. Lancet 2014; 384: 164-72

Neoadjuvant treatment IHC discordancy post treatment

Literature review	Methods	ER discordance	PR discordance	c-erb-2 (Her-2/neu) discordance	Comment
Adams et al. [38]	IHC	2/26 (7.7 %)	4/26 (15.4 %)	6/26 (23.1 %)	Post-NAC on excision
Bogina et al. [8]	IHC	2/36 (5.5 %)	12/36 (33.3 %)		Post-CT and HT on excision
		0/25 (0 %)	2/25 (8.0 %)		Post-CT on excision
		1/24 (4.1 %)	6/24 (25.0 %)		Post-HT on excision
D'Alfonso et al. [39]	IHC/FISH	-	-	14/15 (93.0 %)	Post-NAC on excision
Idinisinghe et al. [12]	IHC	9/49 (18.4 %)	22/41 (53.7 %)	_	LR post-treatment
Kasami et al. [36]	IHC/FISH	19/173 (11.0 %)	27/173 (15.6 %)	Unchanged	Post-NAC on excision
Li et al. [37]	IHC	1.7 % (n = 220)	2.2 % (n = 220)	Unchanged	Post-NAC on excision
Nomura et al. [18]	DCA	7/15 (47 %)	6/6 (100 %)	-	LR post-treatment
Quddus et al. [59]	IHC	-	-	5/39 (12.5 %)	Post-NAC on excision
Rosen et al. [14]	DCA	6/29 (20.7 %)	ND	ND	LR post-treatment

Table 2 Summary of the reported discordant ER, PR, and Her-2/neu cases post-neoadjuvant therapy

ER estrogen receptor, PR progesterone receptor, Her-2/neu epidermal growth factor receptor-2 (c-erb-2), LR local recurrence, NAC neoadjuvant chemotherapy, IHC immunohistochemistry, FISH fluorescent in situ hybridization, DCA dextran-charcoal assay, HT hormone therapy, CT chemotherapy

Breast Cancer Res Treat (2012) 135:29-37

Breast cancer – Molecular intrinsic subtypes

Intrinsic Subtypes Perou et al., Nature 2000 Sorlie et al., PNAS 2001 Sorlie et al., PNAS 2003 Nielsen et al., CCR 2004 Cheang et al., CCR 2008 Parker et al., JCO, Feb 2009 Cheang et al., JNCI 2009 Prat et al., BCR 2010 Nielsen et al., CCR 2010

Breast cancer – Molecular intrinsic subtypes

JOURNAL OF CLINICAL ONCOLOGY

ORIGINAL REPORT

JOURNAL OF CLINICAL ONCOLOGY

EDITORIAL

PAM50 Risk of Recurrence Score Predicts 10-Year Distant Recurrence in a Comprehensive Danish Cohort of Postmenopausal Women Allocated to 5 Years of Endocrine Therapy for Hormone Receptor–Positive Early Breast Cancer

Anne-Vibeke Lænkholm, Maj-Britt Jensen, Jens Ole Eriksen, Birgitte Bruun Rasmussen, Ann S. Knoop, Wesley Buckingham, Sean Ferree, Carl Schaper, Torsten O. Nielsen, Taryn Haffner, Torben Kibøl, Maj-Lis Møller Talman, Anne Marie Bak Jylling, Tomasz Piotr Tabor, and Bent Ejlertsen

Author affiliations and support information (if applicable) appear at the end of this article.

Published at jco.org on January 25, 2018.

Corresponding author: Anne-Vibeke Lænkholm, MD, Department of Surgical

ABSTRACT

Purpose The PAM50-based Prosigna risk of recurrence (ROR) score has been validated in randomized clinical

Do Genomic Assays Provide the Necessary Confidence to De-escalate Adjuvant Therapy?

Ricardo L. B. Costa, H. Lee Moffitt Cancer Center, Tampa, FL William J. Gradishar, Northwestern University, Chicago, IL See accompanying article doi:10.1200/JCO.2017.74.6586

The phrases precision medicine and de-escalation of therapy are being used more frequently in the same sentence when it comes to describing goals of cancer therapy. For perspective, when the National Comprehensive Cancer Network (NCCN) produced its first practice guideline for breast cancer in 1996, the recommendations for adjuvant therapy of early-stage breast cancer were rather simple, reflecting the knowledge generated from clinical trials up to that time.1 Specifically, adjuvant treatment decisions were largely based on age, estrogen receptor (ER) status, tumor size, and the number of axillary nodes involved.

The greater accumulation of clinical trial data married with a far greater understanding of cancer biology has resulted in better outcomes for patients with early-stage disease. Antiestrogen therapy remains the cornerstone of the adjuvant treatment of patients with ER-positive/human epidermal growth factor receptor 2 (HER2)-negative, early-stage breast cancer. Indeed, in a metaanalysis of randomized trials pooling data from 10,645 patients with ER-positive breast cancer, adjuvant treatment with tamoxifen for 5 years significantly reduced not only breast cancer recurrence. rates for 10 years but also led to improvement in the risk breast cancer-related mortality; the relative risk was reduced by approximately 30% throughout the first 15 years from initiation of treatment.2 Adjuvant treatment with chemotherapy can also further reduce the probability of breast recurrence in a subset of patients with localized disease. Results of meta-analyses also conducted under the auspices of the Early Breast Cancer Trialists' Collaborative Group showed that, among 8,575 women, adjuvant treatment with an anthracyclines-based regimen correlated with a relative risk of breast cancer-related mortality of 0.79 when compared with no provided prognostic information and, more importantly, was able to categorize groups of patients with ER-positive, node-negative breast cancer who had such a good prognosis at 10 years with endocrine therapy alone that chemotherapy would not provide additional benefit (predictive).4 The analysis also identified a group at high risk for recurrence at 10 years in whom the added benefit of chemotherapy was clear. There is also an intermediate group in whom the added benefit of chemotherapy was less clear, and it is that subset of patients that is now subject of a large clinical trial (TAILOR-X) to better define the contribution of chemotherapy. The use of this assay has been endorsed by NCCN and ASCO guidelines for over a decade to aid clinical decision-making. The added value of this assay can also be viewed through the lens of clinician recommendations that were changed to, or against, chemotherapy on the basis of results of the assay in patients with node-negative breast cancer.

With an appreciation that it is not clinical features alone but rather the partnering of clinical and molecular features that are the codrivers of any given tumor, the importance of biology has become a key focus in clinical decision-making. For instance, it has long been appreciated that not all node-positive breast cancers will recur even in the absence of any systemic adjuvant therapy. Additionally, even in the era of systemic adjuvant therapy, there are patients with early-stage, ER-positive, node-positive cancer who receive endocrine therapy and in whom disease does not recur in the absence of chemotherapy. Believing that it is more than happenstance and, likely, biology that drives these tumors toward a more favorable clinical course, investigators have explored whether molecular assays may identify those patients with ER-positive.

trials to predict 10-year distant recurrence (DR). The value of Prosigna for predicting DR was examined in a comprehensive nationwide Danish cohort consisting of postmenopausal women with

Major findings from this study – with regards to distant recurrence risk at 10 years after 5 years of endocrine therapy alone

De-escalation of treatment More patients can be spared chemotherapy Immunohistochemical surrogate markers for the molecular intrinsic subtypes

- Limitations
 - No uniform cut off value for Ki67
 - Lack of analytical validity reproducebility
 - Lack of correlation: molecular subtypes and surrogate IHC subtypes

COMMENTARY

Assessment of Ki67 in Breast Cancer: Recommendations from the International Ki67 in Breast Cancer Working Group

Mitch Dowsett, Torsten O. Nielsen, Roger A'Hern, John Bartlett, R. Charles Coombes, Jack Cuzick, Matthew Ellis, N. Lynn Henry, Judith C. Hugh, Tracy Lively, Lisa McShane, Soon Paik, Frederique Penault-Llorca, Ljudmila Prudkin, Meredith Regan, Janine Salter, Christos Sotiriou, Ian E. Smith, Giuseppe Viale, Jo Anne Zujewski, Daniel F. Hayes

Ma

Cor

ortion of cells stween tumor or endocrine

ntre, Fulham Rd,

tak Samuel C Y Leung,¹ Torsten O Nielsen,¹ Lila A Zabaglo,² Ir in 1 Anita L Bane,⁵ John M S Bartlett,^{6,7} Signe Borgquist,⁸ Martin ma Anna Ehinger, ¹¹ Susan Fineberg, ¹² Cornelia M Focke, ¹³ D tex Carolina Gutierrez,15 Judith CHugh,16 Zuzana Kos,17 Annede۱ Mastropasqua,¹⁹ Takuya Moriya,²⁰ Sharon Nofech-Mozes,²¹ Am Penault-Llorca,²² Tammy Piper,⁷ Takashi Sakatani,²³ Rober Tomoharu Sugie,²⁷ Bert van der Vegt,²⁸ Giuseppe Viale,¹⁹ scoring for assessment of Ki67 in breast me ren McShane,³¹ Mitch Dowsett² on behalf of the International Ki bet of the Breast International Group and North American Breas David L. Rimm S, Samuel C. Y. Leung, [...] Mitch Dowsett JΙ

MODERN PATHOLOGY

Article Published: 24 August 2018

An international multicenter study to evaluate reproducibility of automated cancer

Ki67 IHC Identification of hot spots

Poor reproducebility

Immunohistochemical surrogate markers for the molecular intrinsic subtypes

Arch Pathol Lab Med-Vol 140, August 2016

Stains		Luminal BC		Н	HER2 Positive BC			TNBC	
	Luminal A Subtype	Luminal B Subtype (Ki67≥14%)	Luminal B Subtype (PR<20%)	Luminal HER2 PR (≥1%)	Luminal HER2 PR (<1%)	HER2 Enriched	Basal-like subtype	Non- classified subtype	
H&E	C C C	X							
ER	PLE CON								
PR	all's			N	5			State and	
HER2	2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2. 2			10.00					
Ki-67	0000 C		S. S.	C.A.					
CK5	A.A.I			5 4 9 A					
EGFR	040			- Al		1 10 M			

St Gallen international breast cancer conference on primary therapy of early breast cancer – the road of Ki67

Use of pathology to define intrinsic molecular breast cancer subtypes by application of IHC surrogate markers?

2009	Thresholds for therapies. Ki67: 3 categories low <15%, intermediate 16–30% and high >30%
------	--

- 2011 Strategies for breast cancer molecular subtypes genetic testing and attempt for approximation by surrogate IHC markers (ER, PR, HER2 and Ki67) with Ki67 cut off: 14%
- 2013 Personalizing the treatment of women with early breast cancer. Classification of subtypes with Luminal A: ER+, PR ≥20% and Ki67 <20%, HER2-. Luminal B: ER+ and PR<20% and/or Ki67≥20%, HER2-
- 2015 Tailoring therapies-improving the management of early breast cancer: Threshold value of Ki-67 within the range of 20%–29% to distinguish 'luminal B-like` subtype
- 2017 News since St. Gallen 2015: De-escalating and escalating treatment according to stage and breast cancer subtype: "low" ki67 versus "high" ki67
- 2019 Estimating the Benefits of Therapy for Early Stage Breast Cancer: The Panel strongly endorsed the value of genomic assays for determining whether to recommend chemotherapy in T1/T2 N0 tumors, T3 N0 tumors, and TxN1 (1 to 3 positive LN).

Perou CM, Sorlie T, Eisen MB, et al. Molecular portraits of human breast tumours. Nature.2000;406:747–752

Wirapati P et al. Meta-analysis in gene expression profiles in breast cancer: toward a unified understanding of breast cancer subtyping and prognosis signatures. Breast cancer Res 2008; 10: R65

Cheang MCU, Chia SK, Voduc D, et al. Ki67 index, HER2 status, and prognosis of patients with luminal B breast cancer. J Natl Cancer Inst. 2009;101:736–750.

Dowsett M et al. Assessment of Ki67 in breast cancer: recommendations from the International Ki67 in Breast Cancer working group. J Natl Cancer Inst. 2011 Nov 16;103(22)

Lack of correlation: molecular subtypes and surrogate IHC subtype classification

Breast Cancer Res Treat. 2018 Jan;167(1):123-131 DOI 10.1007/s10549-017-4509-9

Digital image analysis outperforms manual biomarker assessment in breast cancer

Gustav Stålhammar^{1,2}, Nelson Fuentes Martinez^{1,3}, Michael Lippert⁴, Nicholas P Tobin⁵, Ida Mølholm^{4,6}, Lorand Kis⁷, Gustaf Rosin¹, Mattias Rantalainen⁸, Lars Pedersen⁴, Jonas Bergh^{1,5,9}, Michael Grunkin⁴ and Johan Hartman^{1,5,7}

Development of an improved panel for basal breast cancer

A survey of immunohistochemical biomarkers for basal-like breast cancer against a gene expression profile gold standard

Jennifer R Won^{1,2}, Dongxia Gao², Christine Chow², Jinjin Cheng², Sherman YH Lau², Matthew J Ellis³, Charles M Perou⁴, Philip S Bernard⁵ and Torsten O Nielsen^{1,2}

¹Department of Pathology and Laboratory Medicine, University of British Columbia, Vancouver, British Columbia, Canada; ²Genetic Pathology Evaluation Centre, University of British Columbia, Vancouver, British Columbia, Canada; ³Division of Oncology, Department of Internal Medicine, Washington University, St Louis, MO, USA; ⁴Department of Genetics, Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA and ⁵Department of Pathology, University of Utah Health Sciences Center, Salt Lake City, UT, USA

- **46** proposed IHC biomarkers published in the literature as associated with the basal subtype
- Utilizing PAM50 gene expression profiling platform as a gold standard

"Nestin positivity or a loss of the expression of inositol polyphosphate-4-phosphate (INPP4B) type 2": the most strongly associated IHC markers with basal like subtype

Sensitivity (83%) and specificity (96%)

Won et al. Mod Pathol. 2013

Scoring of basal markers

Basal-like = Nestin+ OR INPP4B-Non Basal-like = Nestin- AND INPP4B+ ≥1% <5%

Parry et al. J Clin Pathol 2008

Fedele et al. PNAS 2010

Repeat analysis of ER and HER2 in metastatic lesions

Author/ Publication year/ Reference	Number analyzed (ER/HER2/ <i>TOP2A</i>)	Location of biopsy	ER* (%)	HER2* (%)	TOF (%)	°2A*	Comment
Wilking et al (2011) (66)	151	LR+distant ⁵	-	10%	-	No re-a	nalysis ¹
Fabi et al (2011) (67)	137	3/4 LR	-	10%	-		
Amir et al (2010) (51)	258	LR+distant	13%	5%	-	Two pr	ospective studies, pooled
Locatelli et al (2010) (49)	255/167	Distant ⁶	16%	13%	-	No re-a	nalysis ¹
Lindstrom et al (2010) (50)	477/108		33%	10%	-	No re-a	nalysis1,IHC+ICC+biochemical
Karlsson et al (2010) (62)	486	-	35%	-	-	No re-a	nalysis1,IHC+ICC+biochemical
Lower et al (2009) (65)	382	-	-	33%	~	No re-a	nalysis ¹ , IHC only ³
Simmons et al (2009) (54)	25	Distant	12%	8%	-	Prospective study	
Broom et al (2009) (48)	62/18	-	18%	6%	-	No re-a	nalysis'
Liedtke et al (2009) (56)	231	-	18%	14%	-	No re-a	nalysis ¹
Guarneri et al (2008) (55)	75	LR+distant	22%	16%	-	Not all	re-tested ⁴
MacFarlane et al (2008)(180	6) 160	LR+distant	28%	-	-	Total di	iscordance (ER/PgR/HER2)
Tapia et al (2007) (68)	105	Distant ⁶	-	8%	-	IHC (pi	rim BC), ICC (MBC), only FISH
D'Andrea et al (2007) (187)	88/76	syn LN ²	3%	4%	-		
Zidan et al (2005) (64)	58		-	14%	-		
Gong et al (2005) (71)	60	2/3 LR	-	3%	-	1/3 syn	chronous LN, IHC+ICC
Franco et al (2004) (59)	658	-	29%	-	-	A meta	-analysis
Gancberg et al (2002) (69)	93/68	Distant	-	6/7%	-	By IHC	c (6%)/FISH (7%)
Cardoso et al (2001) (188)	370/161	syn LN	-	2%	19%	IHC (T	OP2A, HER2) only
Tanner et al (2001) (70)	46/13	2/3 LR	-	0%	23%	Only TOP2A in 13 pt	
Kuukasjrvi et al (1996) (57)	50	2/3 LR	24%	-	-	Cut-off: ≥ 20 % pos.	

Abbreviations: LN: lymph nodes, LR: locoregional asynchronous disease (i.e. lymph node, scar, and residual breast recurrence), ICC: immunocytochemical analysis."-": No available information. BC: Breast Cancer, MBC: Metastatic Breast Cancer.

^{*}Discordance in percent; ¹No re-analysis done, i.e. based on original pathology reports. ²Assessed on synchronous axillary nodes (i.e. lymph node involvement at diagnosis). ³IHC 2+ scored as HER2 positive. ⁴Did re-evaluate, but not re-test all samples. ⁵The proportion of LR and distant unknown. ⁶Assessed from distant metastases.

- ER discrepancy: 12 29%, often with loss of receptor
- HER2 discrepancy: 6 20%, often with gain of HER2+

Limitations:

- Many "pathology chart review" studies, did not re-analyse tumor samples (methodological variation)
- Prospective studies:

 Treatment decision
 consequence in 15-20%
 Benign disease/other
 malignancies in 14%

Slide courtesy of Jeanette Dupont Jensen. Department of Oncology, Odense University Hospital, Denmark

Mechanism of action of PD-1 and PD-L1 inhibitors

Fig. 1 Mechanism of action of PD-1 and PD-L1 inhibitors. The programmed cell death 1 (PD-1) receptor is expressed on activated T cells, B cells, macrophages, regulatory T cells (Tregs), and natural killer (NK) cells. Binding of PD-1 to its 87 family of ligands, programmed death ligand 1 (PD-L1 or 87-H1) or PD-L2 (87-DC) results in suppression of proliferation and immune response of T cells. Activation of PD-1/PD-L1 signaling serves as a principal mechanism by which tumors evade antigen-specific T-cell immunologic responses. Antibody blockade of PD-1 or PD-L1 reverses this process and enhances antitumor immune activity. TCR, T-cell receptor; MHC, major histocompatibility complex; APC, antigen-presenting cell PD-L1 is expressed on TILs, macrophages, fibroblasts, tumour cells.

Gong et al. Journal for ImmunoTherapy of Cancer (2018) 6:8

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE

Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer

P. Schmid, S. Adams, H.S. Rugo, A. Schneeweiss, C.H. Barrios, H. Iwata, V. Diéras, R. Hegg, S.-A. Im, G.S. Wright, V. Henschel, L. Molinero, S.Y. Chui, R. Funke, A. Husain, E.P. Winer, S. Loi, and L.A. Emens, for the IMpassion130 Investigators* Randomized, prospective, phase 3 trial

– locally adv. or metastastic TNBC
 No prior therapy for advanced TNBC

 Prior chemo in the curative setting, including taxanes, allowed if TFI ≥ 12 mo

Median FU: 12,9 months

Endpoints: PFS, OS

Schmid P. et al. NEJM OCT 2018

IMpassion130 study design

Atezolizumab and Nab-Paclitaxel in advanced Triple-Negative breast cancer phase 3 trial

PD-L1 IHC: Central analysis from archival primary tumor or metastasis **PD-L1 SP142** (Ventana) Cutoff: PD-I $1 \ge 1\%$ IC (immune cells/tumor area) PD-L1 positive N= 369 (40.9%) 185 in atezolizumab group 184 in placebo group

Schmid P. et al. NEJM OCT 2018

Results IMpassion130

		Intent to trea	t	PD-L1 positive (>1%)			
	Tecentriq + Abraxane (n=451)	Placebo + Abraxane (n=451)		Tecentriq + Abraxane (n=185)	Placebo + Abraxane (n=184)		
PFS	7.2	5.5	HR 0.80 (p=0.025)	7.5	5.0	HR 0.62 (p<0.001)	
OS	21.3	17.6	HR 0.84 (p=0.08)	25.0	15.5	HR 0.62*	
*Statistical analysis not carried out owing to failure in ITT population. Source: New England Journal of Medicine.							

Challenges: PD-L1 immunohistochemistry – new biomarker in TNBC

- PD-L1 is a new biomarker for metastatic TNBC in 2019
 - currently only for atezolizumab, but other trials ongoing
- pathologists know PD-L1 from other tumor types (extensive existing training material, currently adapted to TNBC)
- clinicians with a focus on breast cancer will need some basic information to understand the pathology reports
- Typical questions:
 - Which material to apply for analysis? (primary tumour/metastasis)
 - Which antibody to use?
 - Which scoring system?
 - Which cell type?
 - (tumor cell, immune cell (which type of immune cell?)
 - Which cutpoint? depends on clinical setting
 - Reproducebility?

In conclusion

Immunohistochemical classification of breast tumors

- A valuable supplement for the diagnosis of "benign versus in situ" and "in situ versus invasive"
- Histopathological classification of malignant breast tumors
 - Treatment allocation (i.e. lobular vs non lobular)
- Prognostic and predictive factors
 - Selection of treatment and treatment duration
- Intrinsic molecular subtype / gene expression profile
 - Identification of patients who can be spared chemotherapy
- Tumor heterogeneity
 - Repeat analysis
 - multifocal tumors
 - pre/post neo-adjuvant treatment
 - primary tumour/metastasis
- Always keep focus on analytical validity

