

Institute of Pathology, Aalborg University Hospital, Ladegaardsgade 3, P.O.Box 561, DK-9100 Aalborg, Denmark

Recommended protocol for CK8/18

Obtained in run 69

16 Aug 2023

Immunostainer

Type:

Dako Omnis

Primary antibody

Clone:
Producer:
Product no. / lot no.:
Diluent:
Diluent producer / no.:
Dilution factor:
Incubation time / temperature:

5D3 Leica/novocastra NCL-L-5D3 / 6089831 Da Vinci Green **Biocare Medical / PD900M** 1:100 40 min. / 20°C

EnVision Flex / GV800/GV823

Epitope retrieval, HIER

Device:	On Board / On Machine
Buffer:	Dako Omnis Target Retrieval Solution, High pH
Heating time at max. temp .:	30 min.
Maximum heating temp.:	97°C

Dako Omnis

10 min. 20 min. 20°C

Mouse LINKER

Visualization system

Producer:
Product / no:
Linker:
Incubation time linker:
Incubation time polymer:
Incubation temperature:

Chro

hromogen	
Producer:	Dako Omnis
Product / no:	DAB+ Substrate Chromogen System / GV825
Incubation time / temperature:	5 min. / 20°C
Enhancement:	None

Disclaimer:

NordiQC makes every attempt to provide accurate and up-to-date information, yet NordiQC does not make any claim or warranty regarding the accuracy of the provided information nor does it represent that the contents of the web site and protocols reflect the most recent developments in immunohistochemistry at any point in time.